

Analysis on recurrence properties of Weyl's curvature tensor and its Newtonian limit

S. Kumar and K. C. Petwal

Abstract. In this note we present recent enhancements in the cosmological literature related to H.Weyl's conformal curvature tensor. We outline generalized recurrence properties of this tensor in Weyl's manifold and then delineate its Newtonian limit. As well, we discuss some relativistic equations under Newtonian limit, which shall illustrate the role of electricity and magnetism in the geometry of space-time.

M.S.C. 2000: 83F05, 83A05, 53A40.

Key words: Relativistic Weyl space; recurrent Weyl space; Weyl tensor; Newtonian limit; Cotton tensor; Eulerian tensor, Newtonian tidal tensor.

1 Introduction

Hermann Klaus Hugo Weyl (9 November 1885-8 December 1955) is one of the most influential German Mathematicians of 19th century, who dedicated most of his life in developing an enthusiastic geometry now universally known as *geometry of Weyl's space*. Weyl published technical and general works on space, time, matter, philosophy, logic symmetry and history of Mathematics. He was one of the first who visualized general relativity with the laws of electromagnetism. H. Weyl commented that ([17]) "*my work always tried to unite the truth with the beautiful, but when I had to choose one or the other, I usually chose the beautiful*" sums up his efficiency, perfection and curiosity, not only in Mathematics, but in Physics of space-time, matter and Philosophy also. In search of "what is the role of electricity in the geometry of space-time"? Weyl concentrated to this topic in paper after paper and book after book. The standpoints that he and his successors opened up are probably explored today, under the names of "gauge field theory" and "grand unified field theory," by more investigators than there were in the entire physics community at the time of Weyl's first paper in the field. Weyl invented the gauge concept in 1918. By 1928 he had reformulated and restated the idea in the way it is still understood today: "gauge invariance corresponds to the conservation of charge as a coordinate invariance corresponds to the conservation of energy-and-momentum". In a 1918 article, Hermann Weyl ([16, p. 465]) tried to combine electromagnetism and gravity by requiring the theory to be invariant under a local scale change of the metric, i.e., $g_{uv} \rightarrow g_{uv}e^{\alpha(\chi)}$, where χ is a 4-vector.

This attempt was successful and was characterized by Einstein for being inconsistent with observed physical results. It predicted that a vector parallel transported from point p to q would have a length that was path dependent. Similarly, the time interval between ticks of a clock would also depend on the path along which the clock was transported. The article did, however, bears the following crucial facts:

- a) The original term for "Gauge invariance" was "*Erich-invarianz*", which Weyl used in his article. This term refers to invariance under Weyl's scale change.
- b) The geometric interpretation of electromagnetism.
- c) The beginning of non-Abelian Gauge theory. The similarity of Weyl's theory to non-Abelian Gauge theory is more striking in his 1929 paper¹.

Now, in order to pursue our proposed study "A review on recurrence properties of Weyl's curvature tensor and its Newtonian limit", we briefly introduce some notions on Weyl's space, Generalized Weyl's space and Newtonian limit of general relativity.

Weyl's space

An n -dimensional differentiable manifold W_n is said to a Weyl space, if it has a symmetric connection ∇ and a symmetric conformal metric tensor g_{ij} preserved by ∇ satisfying the compatibility condition given by the equation ([10],[12],[15])

$$(1.1) \quad \nabla_k g_{ij} - 2T_k g_{ij} = 0,$$

where T_k represents a covariant vector field.

Re-writing (1.1) in an extended fashion, we have

$$(1.2) \quad \frac{\partial}{\partial x^k} g_{ij} - g_{hj} \Gamma_{jk}^h - 2T_k g_{ij} = 0,$$

where Γ_{kl}^i are the connection coefficients of the symmetric connection ∇ and are defined as

$$(1.3) \quad \Gamma_{kl}^i = \binom{i}{k \ l} - g^{im} (g_{mk} T_l + g_{ml} T_k - g_{kl} T_m).$$

Moreover, under the renormalization condition:

$$(1.4) \quad \tilde{g}_{ij} = \lambda^2 g_{ij},$$

of the metric tensor g_{ij} , the covariant vector field T_k is transformed by the law:

$$(1.5) \quad \tilde{T}_k = T_k + \frac{\partial}{\partial x^k} \ln \lambda,$$

where λ is a scalar function defined on W_n . Thus the space W_n satisfying all the foregoing condition will be symbolized by $W_n(\Gamma_{jk}^i, g_{ij}, T_k)$, or $W_n(g, T)$. Also, a geometric object Ω defined on $W_n(\Gamma_{jk}^i, g_{ij}, T_k)$ is called a satellite of weight $\{w\}$, of a tensor g_{ij} , if it admits a transformation of the form:

$$(1.6) \quad \tilde{\Omega} = \lambda^w \Omega$$

¹Weyl, H., Proc. Natl. Acad. Sci. 15 (1929), 32.

under renormalization condition of the metric tensor g_{ij} ([10],[5]). Further, the prolonged covariant derivative of a satellite Ω is defined by:

$$(1.7) \quad \dot{\nabla}_k \Omega = \nabla_k \Omega - w T_k \Omega.$$

It is remarkable that the prolonged derivative preserves the weight.

Generalized Weyl's space

An n -dimensional differentiable manifold GW_n having an anti-symmetric connection ∇^* and anti-symmetric metric tensor g_{ij}^* , preserved by ∇^* is called a "generalized Weyl space" [9]. For such a space, in local co-ordinate system, we have a compatibility condition as below:

$$(1.8) \quad \nabla_k^* g_{ij}^* - 2T_k^* g_{ij}^* = 0,$$

where T_k^* are the components of a covariant vector field called the complementary vector field of the GW_n space. Using the concept of covariant differentiation ([18],[8]), the compatibility condition, (1.8) can be written in extended form as:

$$(1.9) \quad \frac{\partial^*}{\partial x^k} g_{ij}^* - g_{hj}^* L_{ik}^h - g_{ih}^* L_{jk}^h - 2T_k^* g_{ij}^* = 0,$$

where L_{kl}^i are the connection coefficients of the anti-symmetric connection ∇_k^* and are obtained from the compatibility condition as [1]:

$$(1.10) \quad L_{kl}^i = \Gamma_{kl}^i + \frac{1}{2} [\chi_{km}^h g_{lh}^* + \chi_{ml}^h g_{kh}^* + \chi_{kl}^h g_{hm}^*] g^{*mi}.$$

Now, putting

$$(1.11) \quad \chi_{kl}^i = \frac{1}{2} [\chi_{km}^h g_{lh}^* + \chi_{ml}^h g_{kh}^* + \chi_{kl}^h g_{hm}^*] g^{*mi},$$

we obtain

$$(1.12) \quad L_{kl}^i = \Gamma_{kl}^i + \chi_{kl}^i,$$

where Γ_{kl}^i and χ_{kl}^i are respectively the coefficients of a Weyl connection and the torsion tensor of GW_n space and are expressed as:

$$(1.13) \quad \Gamma_{kl}^i = \frac{1}{2} (L_{kl}^i + L_{lk}^i) = L_{(kl)}^i,$$

(here, the round bracket stands for symmetry.) and

$$(1.14) \quad \chi_{kl}^i = \frac{1}{2} (L_{kl}^i - L_{lk}^i) = L_{[kl]}^i$$

(here the square bracket stands for anti-symmetry).

A generalized Weyl space satisfying all the aforementioned conditions is symbolized as $GW_n (L_{jk}^i, g_{ij}^*, T_k^*)$.

Newtonian limit in General Relativity

Some researchers attempted to establish a closed Newtonian system for the evolution of fluid quantities (for instance, density, expansion, shear etc.) by considering Newtonian limit of the corresponding equation in general relativity. Afterward, a considerable amount of research has been dedicated towards the supporting or disproving this goal by [6] and many other researchers. Actually, the concept of Newtonian limit in theory of relativity has been introduced to focus on two major aspects; one of them concerns to the presentation of a precise derivation of Newtonian limit of fluid evolution equation in a 4-dimensional "frame theory" developed by ([3], [2]). This theory covers both Einstein and Newtonian theory of gravitationally interacting matter. On the other hand, second aspect pertains to the discussion general relativistic equations to describe a closed Newtonian system.

2 Recurrence properties of Weyl's curvature tensor in W_n and GW_n spaces

According to [10], under a renormalization condition of the fundamental metric tensor g_{ij} of the form (1.4), an object Ω defined on GW_n space, admits a transformation of the form (1.6) is called a satellite with weight $\{w\}$ of the metric tensor and the prolonged covariant derivative of the satellite Ω relative to the symmetric connection ∇ is defined by [11]:

$$(2.1) \quad \dot{\nabla}_k \Omega = \nabla_k \Omega - w T_k \Omega.$$

Whereas, the same relative to anti-symmetric connection ∇^* is defined as:

$$(2.2) \quad \dot{\nabla}_k^* \Omega = \nabla_k^* \Omega - w T_k^* \Omega,$$

which evinces that the prolonged derivative preserves the weight of satellite.

Now, the conformal Weyl curvature tensor C_{ijkl}^i of the generalized Weyl space GW_n is given by [7] as:

$$(2.3) \quad C_{jkl}^i = R_{jkl}^i + \frac{2}{n(n-2)} (\delta_l^i R_{[jk]} - \delta_k^i R_{[jl]} - g_{jl} g^{im} R_{[mk]} + g_{jk} g^{im} R_{[ml]} - (n-2) \delta_j^i R_{[lk]}) - \frac{1}{(n-2)} (\delta_l^i R_{jk} - \delta_k^i R_{jl} - g_{jl} g^{im} R_{mk} + g_{jk} g^{im} R_{ml}) + \frac{R}{(n-1)(n-2)} (g_{jk} \delta_l^i - g_{jl} \delta_k^i),$$

where the square bracket stands for the anti-symmetrization.

This n -dimensional GW_n space is said to be conformally recurrent Weyl space if its conformal curvature tensor (2.3) of weight $\{0\}$, on taking the prolonged derivative, satisfies the condition:

$$(2.4) \quad \dot{\nabla}_m^* C_{jkl}^i = \phi_m C_{jkl}^i,$$

where $\phi_m \neq T_m$ is a non-zero covariant recurrence vector field of weight $\{0\}$.

The conformal curvature tensor given by (2.3) can be re-defined in the purely contravariant pattern as below:

$$(2.5) \quad C^{ijkl} \equiv R^{ijkl} - \frac{1}{(n-2)} (g^{il} S^{kj} - g^{il} S^{jk} - g^{jk} S^{li} + g^{jl} S^{ki}),$$

where $S^{lj} \equiv R^{lj} - \frac{1}{2(n-1)}g^{lj}R$. R^{lj} and R are the Ricci quantities, while C^{ijkl} bears the following properties:

(a). It is purely covariant against conformal redefinition of the metric, i.e., $g_{kl}(x) \rightarrow \lambda(x)g_{kl}(x)$.

(b). It vanishes if and only if the GW_n space is conformally flat, i.e., g_{kl} is diffeomorphic to $\lambda\eta_{kl}$, where η_{kl} is flat.

(c). It possesses the symmetries of Riemannian tensor and also is traceless in each index pair.

(d). Evidently, from the properties (a) and (b), the Weyl tensor acts as a template for conformal flatness. Thus by evaluating it on a specific metric tensor, one can distinguish, whether the space-time under consideration is conformally flat or not.

In three dimensions the Weyl tensor vanishes identically and the Riemannian tensor is given by the last term in (2.5) at ($n = 3$). But not all three dimensional space-times are conformally flat.

Now, as it is well known that in dimensions greater than three, the conformal tensor (2.3) and (2.5) is the Weyl tensor, then what about the three dimensional space-time? To overcome from this difficulty, one need a substitute for Weyl tensor, which would act as a template for the conformal flatness. Indeed, there is a crucial substitution for Weyl tensor, known as *Cotton Tensor*, which is delineated as:

$$(2.6) \quad C^{ij} = \frac{1}{2\sqrt{g}} \left(\epsilon^{ikl} D_k R_l^j + \epsilon^{jkl} D_k R_l^i \right).$$

This Cotton tensor serves that role, as it possesses conformal template properties (a) and (b). C^{ij} is symmetric in its indices and like the Weyl tensor, it is traceless.

Therefore, in case of the space-time having dimension less than 4, one can have the conformal recurrence properties for such conformally non-flat continuum by taking the prolonged derivative of Cotton tensor (2.6) of weight $\{0\}$. The conformally non-flat space-time having dimension $n = 3$ will be called conformally recurrent if the Cotton tensor satisfy the following:

$$(2.7) \quad \dot{\nabla}_m^* C^{ij} = \phi_m C^{ij},$$

where ϕ_m is a non-zero recurrence vector field of weight zero.

It is remarkable that for the space-time having dimension less than three, there is no need of a conformal tensor and indeed none exists as all such spaces are locally conformally flat.

We, now, study conformal Weyl tensor, its Newtonian limit and some relativistic equations in general relativity due to Newtonian limit.

3 Conformal Weyl's curvature tensor, its Newtonian limit and relativistic equations

In the modern cosmological literature, role of conformal Weyl's curvature tensor has been adequately discussed. Especially, concentration upon the crucial parts, namely "electric" and "magnetic" parts of conformal Weyl's curvature tensor has been drawn by ([3], [2]) individually. Further, ([3], [2]) have discussed some ideas on the magnetic part of Weyl tensor under Newtonian limit of general relativity. In order to pursue

significance of Weyl tensor in cosmological structures, the "frame-theory" for a general 4-dimensional space-time continuum is employed, which encapsulate both the "Newtonian's theory" as well as "Einstein's theory". In this frame theory, a single parameter $\varepsilon = c^2$ is introduced in such a way that it distinguishes between Newtonian's and Einstein's theories. The limit for this parameter is taken as $\varepsilon \rightarrow 0$. Moreover, in the formalism of Einstein's theory, a temporal metric t_{ij} and an inverse spatial metric s^{ij} are used which are related by an expression of the form:

$$(3.1) \quad t_{ij}S^{jk} = -\varepsilon\delta_i^k.$$

In case of general relativity, the parameter ε is taken to be greater than zero, i.e., $\varepsilon > 0$ and the Riemannian as well as inverse Riemannian metric g_{ij} and g^{ij} are used such that:

$$(3.2) \quad g_{ij} = -\varepsilon^{-1}t_{ij} \text{ and } g^{ij} = S^{ij}.$$

On the other hand, in Newtonian's theory, the parameter ε is taken to be zero and the temporal metric is defined as:

$$(3.3) \quad t_{ij} = t_{,i}t_{,j},$$

where t is the absolute time and the comma (,) denotes the derivative with respect to the Eulerian co-ordinates, which will be discussed further.

In the conception of frame theory, the conformal Weyl's curvature tensor (2.3), for $\varepsilon > 0$ takes the following form:

$$(3.4) \quad C_{jkl}^i = R_{jkl}^i - \delta_k^i R_{lj} - \varepsilon^{-1} \left[t_{jk} R_{lm} s^{mi} + \frac{1}{3} \delta_k^i t_{lj} R_{pr} z^{pr} \right].$$

This expression becomes worthless for $\varepsilon = 0$. However, if one uses the Einstein's field equation:

$$(3.5) \quad R_{ij} = 8\pi G \left(t_{ik} t_{jl} - \frac{1}{2} t_{ij} t_{kl} \right) T^{kl} - \Lambda t_{ij}$$

of the frame theory (valid for $\varepsilon \geq 0$), to eliminate R_{ij} from the equation (3.4), we obtain:

$$(3.6) \quad C_{jkl}^i = R_{jkl}^i - 8\pi G \left[\delta_k^i t_{lp} t_{jr} T^{pr} - t_{jk} t_{lm} T^{mi} - \frac{2}{3} \delta_k^i t_{lj} t_{pr} T^{pr} \right].$$

Now, this formula is noteworthy, even for $\varepsilon = 0$. Thereby, one can define the conformal Weyl's curvature tensor in the frame theory by equation (3.6). This expression for the Weyl's curvature tensor is quite suitable, for instance, if a sequence of general relativistic solutions has a Newtonian solution as a limit, then the limit of conformal Weyl curvature tensor is surely produced by (3.6). Furthermore, in case of Newtonian's theory, ($\varepsilon = 0$), equation (3.6) due to equation (3.3) reduces to the form:

$$(3.7) \quad C_{jkl}^i = R_{jkl}^i - \frac{8\pi G}{3} \rho t_{,j} \delta_k^i t_{,l}.$$

Also, the "electric" and "magnetic" part of the conformal Weyl curvature tensor with respect to any 4-velocity vector v^i can be obtained from (3.7) and these are respectively defined like below:

$$(3.8) \quad E_k^i = R_{kjl}^i v^j v^l - \frac{4\pi G}{3} \rho (\delta_k^i - v^i t_{,k});$$

$$(3.9) \quad H_{ik} = \frac{1}{2} \eta_{ijpr} s^{ru} C_{ukl}^p v^j v^l = 0.$$

Here it is very observable that the magnetic part H_{ik} of the conformal Weyl tensor vanishes in the Newtonian limit. This fact can be more precisely justified as follows:

In the general relativity, H_{ik} measures the relative rotation of nearby freely falling gyroscopes due to gravitomagnetism. This effect has nonexistence in case of Newtonian theory in which the parallelism of spatial vectors is path independent. It means, parallel gyroscopes will always remain parallel if subjected to nothing except inertia and gravity.

Here, we now discuss some relativistic equations in Newtonian limit.

In three dimensional Weyl space, the equation (3.8) yields a new kind of tensorial quantity called "Newtonian tidal tensor" which is trace free part of gravitational field tensor $g_{(i,j)}$ (here comma denotes derivative with respect to Eulerian co-ordinate system) and is given by

$$(3.10) \quad E_{ij} = g_{i,j} - \frac{1}{3} \delta_{ij} g_{l,l},$$

provided

$$(3.11) \quad E_{[ji]} = 0; E_{ii} = 0.$$

Likewise any Eulerian field, the Newtonian tidal tensor of the gravitational field strength \vec{g} can be written in terms of Lagrangian co-ordinates as below:

$$(3.12) \quad E_{ij} = g_{i|k} J_{kj}^{-1} - \frac{1}{3} \delta_{ij} g_{l|k} J_{kl}^{-1},$$

where a vertical slash stands for the derivative with regards to Lagrangian co-ordinate system.

In order to discuss evolution of tidal tensor as relativistic equation, we introduce a diffeomorphism $\vec{f}_t : \vec{x} = \vec{f}(\vec{X}, t)$, which sends fluid elements from their initial

Lagrangian position \vec{X} to a point \vec{x} in the Eulerian space at time t . Also, we use an expression for the Jacobian of the inverse transformation:

$$(3.13) \quad \vec{h} = \vec{f}^{-1}, \vec{g}[\vec{x}, t] = \vec{\ddot{f}}\left(\vec{h}[\vec{x}, t], t\right), J = \det(f_{i|k}),$$

and

$$(3.14) \quad h_{j|l} = \frac{1}{2J} \varepsilon_{jprq} \tau(\vec{\ddot{f}}_l, f_p, f_q) - \frac{1}{3} \varepsilon_{opq\tau}(\vec{\ddot{f}}_o, f_p, f_q) \delta_{ij},$$

so that the Newtonian tidal tensor could explicitly be expressed in terms of \vec{f} as:

$$(3.15) \quad E_{ij} = \frac{1}{2J} \left[\varepsilon_{jprq} \tau(\ddot{f}_l, f_p, f_q) - \frac{1}{3} \varepsilon_{opq} \tau(\ddot{f}_o, f_p, f_q) \delta_{ij} \right].$$

Therefore, any trajectory field \vec{f} , which obeys the Lagrange-Newtonian system is given as:

$$(3.16) \quad \tau(\ddot{f}_J, f_j, f_k) = 0,$$

and

$$(3.17) \quad \tau(\ddot{f}_1, f_2, f_3) + \tau(\ddot{f}_2, f_3, f_1) + \tau(\ddot{f}_3, f_1, f_2) - \Lambda J = -4\pi G \overset{0}{\rho},$$

where Λ is a cosmological constant.

The last two equations determine the evolution of tidal tensor in the form of relativistic equation of general relativity via equation (3.12). In equations (3.16) and (3.17), the symbol $\tau(A, B, C)$ etc. denote the functional determinant of any functions $A(\vec{X}, t)$, $B(\vec{X}, t)$ and $C(\vec{X}, t)$ with respect to Lagrangian co-ordinate system (\vec{X}) and $\overset{0}{\rho}$ denotes the initial density field [4].

4 Concluding remarks

In the present manuscript, we have tried to draw our focus on recurrence properties of conformal Weyl's curvature tensor and its applications in the modern literature of relativity and cosmology. Particularly, we have discussed Weyl space, generalized Weyl space and their conformal curvature tensors along with recurrent nature. It is shown that for the space-time having dimensions less than 4, needed a tensor (called Cotton tensor), other than Weyl's tensor to check out the conformal flatness of the space-time and its recurrent nature. Moreover, a relativistic form of Weyl's tensor and relativistic equation evolved due to its parts (namely, electric and magnetic) has been studied.

Acknowledgment. The authors wish to acknowledge the funding agency as the present research work is financially assisted by the organization "U-COST", Dehradun. Authors also wish to extend their immense thanks to all those scientists/researchers, whose articles have been used during the preparation of this article. Especially, hearty thanks are also due to Professors J. Ehlers, T. Buchert and L.Z. Akgün as their original epics have been used herein.

References

- [1] G. Civi and A. Özdeger, *Theory of Nets in generalized Weyl spaces*, Webs and Quasigroups, Tver State Univ., in print.
- [2] J. Ehlers, *The Newtonian limit of General Relativity*, In (Ed.: G. Fenarese) "Classical Mechanics and Relativity, Relationship and Consistency", Naples, Bibliopolis, 1991, 95-106.

- [3] J. Ehlers, *Über den Newtonschen Grenzwert der Einsteinschen Gravitationstheorie in Grundlagenprobleme der Modernen Physik* (in German), In (Eds.: J. Nitsch, J. Pfarr, E.W. Stachow): "Grundlagenprobleme der Modernen Physik", Bibliograph. Inst. Mannheim, 1981, 65-84.
- [4] J. Ehlers and T. Buchert, *On the Newtonian limit of the Weyl tensor*, arXiv: 0907.264v1 [gr-qc], (2009), 1-6.
- [5] V. Hlavaty, *Theorie d'immersion d'une W_m dans W_n* , Ann. Soc. Polon. Math. 21 (1949), 196-206.
- [6] S. Matarrese, *Course CXXXII "Dark Matter in The Universe"*, Varenna, Proc. IOP 'Enrico Fermi', 1995.
- [7] R. Miron, *Mouvements confirmés dans les espaces W_n et N_n* , Tensor N.S. 19 (1968), 33-41.
- [8] V. Murgescu, *Espaces de Weyl generalises*, Bul. Inst. Pol. de Jassy, 1970.
- [9] V. Murgescu, *Espaces de Weyl generalises*, Rev. Roum. Math. Pures et Appl. XV, (2) (1970), 293-301.
- [10] A. Norden, *Affinely connected spaces*, GRMFL Moscow, (1976).
- [11] A. Norden and S. Yafarov, *Theory of non-geodesic vector field in two dimensional affinely connected spaces*, Izv. Vuzov, Matem., 12 (1974), 29-34.
- [12] B. Tsareva and G. Zlatanov, *On the geometry of the nets in the n-dimensional space of Weyl*, J. Geom. 38 (1990), 182-197.
- [13] C. Udriste, V. Balan and A. Udriste, *Magnetic fields generated by piecewise rectilinear configurations*, Appl. Sci., 1, 1 (1999), 12-39.
- [14] C. Udriste, M. Postolache, A. Soeanu, *Computer simulation of magnetic phase portraits and geometric dynamics around piecewise rectilinear circuits*, Appl. Sci., 1, 1 (1999), 40-49.
- [15] G. Zlatanov, *Nets in the n-dimensional space of Weyl*, C.R. Acad. Bulgari Sci. 41 (1998), 29-32.
- [16] H. Weyl, *Sitzungsber. Preuss, Akad. Wiss. Berlin (Math. Phys.)* 1918, 465.
- [17] J.A. Wheeler, *Hermann Weyl and unity of knowledge*, American Scientist 74 (1986), 366-375.
- [18] L. Zeren Akgun, *On generalized Weyl spaces*, Bull. Calcutta Math. Soc. 91, (4) (1999), 267-278.

Authors' address:

S. Kumar and K.C. Petwal
 Department of Mathematics, HNB Garhwal University,
 Campus Badshahi Thaul, Tehri Garhwal, Uttarakhand, 249 199, India.
 E-Mail: sandeep_2297@rediffmail.com, drsandeepbahuguna@rediffmail.com,
 drkcpetwal@rediffmail.com